




EFFECTS ON COTTON YIELDS

California – 2008



Plot Information:

Cooperator: Bill Weir, Ph.D. & Pafford Farms, Dos Palos, CA
Variety: Phytogen 725
Planted: April 7, 2008
Replications: 4
Treatments: 4 rows x 30 inch beds x 50 ft in length.

Harvested: Oct. 9, 2008
Plot Design: RCBD

Fertility:

Prior to planting, the soil received 230 #/A Ammonium sulfate, and 40 gal/A, 4-10-10 – Zn.


Application Information:

Products were applied to the tops of the beds on either side of the drill row, and worked in using hand tools on April 17, 2008.



EFFECTS ON COTTON YIELDS

California – 2008



Plot Information:

Cooperator: Bill Weir, Ph.D. & Pafford Farms, Dos Palos, CA
Variety: Phytogen 725
Planted: April 7, 2008
Replications: 4
Treatments: 4 rows x 30 inch beds x 50 ft in length.

Harvested: Oct. 9, 2008
Plot Design: RCBD

Fertility:

Prior to planting, the soil received 230 #/A Ammonium sulfate, and 40 gal/A, 4-10-10 – Zn.

Application Information:

Products were applied to the tops of the beds on either side of the drill row, and worked in using hand tools on April 17, 2008.